

Spectral properties of 3EG gamma-ray sources outside the galactic plane

O. Reimer (MPE Garching, D-85740 Garching, Germany), D.L. Bertsch, R.C.Hartman, S.D. Hunter, D.J. Thompson (Code 661, NASA GSFC, Greenbelt, MD 20771, USA), P.F. Michelson, Y.C. Lin, P.L. Nolan (W.W.Hansen Experimental Physics Lab, Stanford University, Stanford, CA 94305, USA), G. Kanbach, H.A. Mayer-Hasselwander (MPE Garching, D-85740 Garching, Germany), D.A. Kniffen (Hampden-Sydney College, Dept. of Physicsm Hampden-Sydney, VA 23943), R. Mukherjee (Barnard College, Dept. of Physics, New York, NY 10027, USA), B.L. Dingus (Physics Dept., University of Utah, Salt Lake City, UT 84112), P. Sreekumar (HEAD, ISRO Satellite Center, Bangalore, India)

Most of the high-energy gamma-ray sources within the third EGRET catalogue are accompanied by their spectral index. However, a more closer look revealed, that in various cases single power-law fits show significant difficulties to explain the data between 30 MeV and 10 GeV in a sufficient way. Additionally, deviations from a common spectral index through all available EGRET observations are already reported. This included currently a few AGN, like 3C279 and 1622-297, in different states of activity. Here a detailed spectral study for the EGRET sources outside the galactic plane is presented. In particular, spectral details on unidentified high-latitude sources are provided. Deviations from simple single power-law fits are investigated, and alternative fits are suggested in appropriate cases.